



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search PubMed  for  Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract  Sort  Save Text Clip Add Order

1: J Chromatogr A 1998 Dec 11;827(2):337-44

Related Articles, Books, LinkOut

## New approaches to the isolation of DNA by ion-exchange chromatography.

Levison PR, Badger SE, Hathi P, Davies MJ, Bruce IJ, Grimm V.

Whatman International Ltd., Maidstone, Kent, UK.

The performance of different anion-exchange media have been compared for the isolation of plasmid DNA and genomic DNA from bacterial cells and human whole blood. Whatman DEAE-Magarose, based on an agarose bead containing a paramagnetic component, has been compared with prepacked gravity-flow columns containing a derivatised silica matrix. In each case the DNA isolation at various scales of operation was similar both in terms of yield and quality. The magnetic susceptibility of DEAE-Magarose is very high, facilitating the use of this separation technique for rapid flexible batch chromatographic processes, a limitation of the prepacked column techniques.

PMID: 9914660 [PubMed - indexed for MEDLINE]

Display Abstract  Sort  Save Text Clip Add Order

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
Department of Health & Human Services  
[Freedom of Information Act](#) | [Disclaimer](#)